

Idiopathic Thrombocytopenic Purpura in Two Mothers of Children With DiGeorge Sequence: A New Component Manifestation of Deletion 22q11?

A. Lévy,^{1*} G. Michel,² M. Lemerrer,³ and N. Philip¹

¹Department of Genetics and INSERM U406, Hopital d'Enfants de la Timone, Marseilles, France

²Department of Paediatric Hematology, Hopital d'Enfants de la Timone, Marseilles, France

³Department of Genetics and INSERM U393, Hopital des Enfants Malades, Paris, France

The phenotypic spectrum caused by the microdeletion of chromosome 22q11 region is known to be variable. Nearly all patients with DiGeorge sequence (DGS) and approximately 60% of patients with velocardiofacial syndrome exhibit the deletion. Recent papers have reported various congenital defects in patients with 22q11 deletions. Conversely, some patients have minimal clinical expression. Ten to 25% of parents of patients with DGS exhibit the deletion and are nearly asymptomatic. Two female patients carrying a 22q11 microdeletion and presenting with idiopathic thrombocytopenic purpura are reported. Both had children with typical manifestations of DGS. Am. J. Med. Genet. 69:356–359, 1997. © 1997 Wiley-Liss, Inc.

KEY WORDS: DiGeorge sequence; deletion 22q11; idiopathic thrombocytopenic purpura; immune dysfunction

INTRODUCTION

The phenotypic expression of the microdeletion of chromosome 22 within the 22q11 region is known to be highly variable [Driscoll et al., 1993; Wilson et al., 1993; Levy-Mozziconnacci et al., 1994]. Most patients reported to date have the DiGeorge sequence (DGS) or velocardiofacial syndrome (VCFS). These two conditions have been described separately and are clinically different but with some degree of overlap. Patients with milder clinical pictures, either isolated congenital conotruncal cardiac defects [Goldmuntz et al., 1993; D'Angelo et al., 1994] or hypoparathyroidism [Sciré

et al., 1994] have been reported. In 10 to 25% of cases, one of the parents is found to be a carrier of the deletion [Wilson et al., 1993]. Most of these parents are nearly asymptomatic. We report on two families in which the mothers exhibit the deletion and have idiopathic thrombocytopenic purpura (ITP).

CLINICAL REPORTS

Family 1

The mother was born in 1967. She had learning disability and speech difficulties, with hypernasal voice. Physical examination showed large nose, short and poorly defined philtrum, and small ears with an overfolded helix (Figs. 1 and 2).

At age 20, she developed severe thrombocytopenia with less than 10,000 platelets per mm³. ITP was diagnosed on the following criteria: spleen and liver were not enlarged, the megakaryocyte number was increased in bone marrow smears with apparently normal granulopoiesis and erythropoiesis, HIV serology was negative, the sedimentation rate was normal, and there were no antinuclear antibodies. Isotopic study of platelet survival with ⁵¹Cr radiolabelled alloplatelets showed a markedly decreased lifespan (less than 1 day) and an almost exclusive splenic sequestration. Interestingly, this woman also had chronic leuconutropenia with WBC counts ranging from 1,900 to 4,000/mm³ on different examinations, and neutrophil counts between 500 and 2300/mm³ and lymphocyte counts between 900 and 2,300/mm³ (Table I). The patient received several courses of corticotherapy and intravenous immunoglobulin, which only resulted in transient improvements of the thrombocytopenia. Finally, she underwent splenectomy and had complete recovery of her platelet counts. Two years after splenectomy, the last blood count showed 187,000 platelets. The mean platelet volume was slightly increased (11.5 µm³) but there were no giant platelets. A flow cytometry study of platelet glycoproteins, using monoclonal antibodies specific for GPIb, GPIIb/GPIIIa, and GPIa/IIa, showed a normal expression of these glycoproteins. The lymphocyte count was 2,480/mm³ with 1,580 CD3+ cells/mm³.

Contract grant sponsor: Groupement de Recherches et d'Etudes sur les Génomes; Contract grant sponsor: EEC.

*Correspondence to: A. Lévy, Centre de Genetique Medicale Hôpital d'Enfants de La Timone 13385, Marseille Cedex 5, France.

Received 19 June 1995; Accepted 20 September 1995



Fig. 1, 2. Frontal and profile view of the mother of family 1, showing bulbous nose, short philtrum, and small ears.

In 1993, she became pregnant. Pregnancy and delivery were uneventful. At birth, the child had heart murmur. Interrupted aortic arch and VSD were diagnosed. The boy had minor facial anomalies, with round face, upslanted palpebral fissures, hypertelorism, small mouth, and small and malformed ears (Fig. 3). Transient hypocalcemia was documented. The congenital heart defect was repaired at age 1 month. DGS was suspected.

Fluorescent in situ hybridization (FISH) with cosmid probes sc4.1 and sc11.1, specific of the 22q11 region, was performed according to the methods previously described [Levy-Mozziconacci et al., 1994]. Cosmid probes were labelled by nick-translation with biotin 11-dUTP, according to BRL protocol. The hybridization signal was evoked by FITC labelled-avidin, amplified twice with additional layers of biotinylated goat anti-avidin and avidin-FITC, and chromosomal DNA was counterstained by propidium iodide. The mother and child were deleted for both loci.

Family 2

In this family, the first child, born in 1991, had DGS with interrupted aortic arch, VSD, hypocalcemia, and absent thymus. No molecular studies were conducted. This child died at age 1 month and the couple was referred for genetic counseling. The 25-year-old mother was found to have a long nose and a nasal speech

(Fig. 4). Personal history documented developmental delay. She walked at age 4 years.

A low platelet count was detected at age 26 years, during her second pregnancy. Thrombocytopenia was moderate (platelet counts between 90,000 and 120,000/mm³) but chronic, now persisting for 3 years after the initial evaluation. There was no hepatosplenomegaly and antinuclear antibodies were negative. Considering these facts, the woman was thought to have mild ITP which did not require any treatment.

The child, a girl, was born by C-section. She exhibited severe neonatal hypocalcemia with seizures. There was no visible thymic shadow on X-ray films. Facial appearance was typical of DGS. Transient thrombocytopenia was observed during the first days of life, as is usual in neonates born to mothers with ITP. This thrombocytopenia was successfully treated with intravenous immunoglobulins. FISH analysis of the 22q11 region demonstrated deletion in both mother and child.

DISCUSSION

The clinical spectrum of microdeletion 22q11 has widened. It is now recognized that such a deletion can lead to a variety of phenotypes. DGS, characterized by thymic hypoplasia, hypocalcemia, congenital heart defect of conotruncal type, and typical facies, represents the more severe phenotype [Wilson et al., 1993]. VCFS,

TABLE I. Hematological Findings

	Mother of family 1			Mother of family 2
	Before splenectomy		After splenectomy	
Platelet count ^a	8,000	11,000	187,000	120,000
WBC count ^a	2,400	1,900	3,400	6,000
Neutrophil count ^a	U*	800	310	4,740
Lymphocyte count ^a	U	900	2,480	1,020
Monocyte count ^a	U	150	540	250
CD3+ lymphocytes ^a	U	U	1,580	U

* U, unknown.

^a Per mm³.



Fig. 3. Frontal view of the child of family 1. Note the small mouth.

described in 1978 by Shprintzen et al. in patients with velopharyngeal insufficiency, congenital heart defect, and learning disabilities is also found to be associated with 22q11 deletion in most cases. Wilson et al. [1993] proposed the acronym CATCH 22 (Cardiac defects, Abnormal facies, Thymic hypoplasia, Cleft palate, Hypocalcemia) to describe the clinical spectrum of microdeletion 22q11. However, these five letters are far from describing all the component manifestations that can be found in patients hemizygous for 22q11. A variety of other congenital defects have been reported in patients with DGS [Wilson et al., 1993; Nickel et al., 1994]. Studies of patients with VCFS have shown that late-onset disorders, such as psychosis, can complicate

the prognosis of these children. Conversely, very mild forms have been reported [Lipson et al., 1994], including 10 to 25% of parents of children with DGS, who exhibit the molecular defect and represent the milder end of the phenotype.

We present two cases of mothers whose 22q11 hemizygosity was detected after the birth of a child with fully symptomatic DGS. Both mothers had a mild form of VCFS and both had significant and prolonged low platelet counts. Thrombocytopenia is not a classical manifestation of either DGS or VDFS. Recently, Budarf et al. [1994] reported on a case of 22q11 deletion associated with Bernard-Soulier syndrome. This rare congenital bleeding disorder, characterized by thrombocytopenia and very large platelets, is associated with a defect in glycoprotein 1b (GP1b), the major platelet receptor for von Willebrand factor. As the Gp1b β has been localized within the DG critical region, the authors postulated that the haploinsufficiency of 22q11 in this patient unmasked a mutation in the other allele. In our first patient, this condition can be clearly excluded because there were no giant platelets and because GP1b was normally expressed on the platelet membrane. In both patients, the clinical course suggests ITP. Two cases of DGS with ITP have been reported so far [Jyonouchi et al., 1991]. Of course, this occurrence could be coincidental. However, defective thymic function may predispose patients with DGS to autoimmune diseases. The two cases reported here suggest that patients with minimal clinical expression, such as parents of children with typical DGS, are also predisposed to ITP and that these ITP might be due to subtle dysfunction of T-cell regulation. Moreover, these two observations underline that ITP can be the first apparent pathologic event in the life of patients with the 22q11 microdeletion.

Careful examination of the clinical history of parents carrying the 22q11 microdeletion will certainly lead to a better definition of the long-term prognosis of deletion 22q11.

ACKNOWLEDGMENTS

We thank Professor JF Schved for providing clinical information on patient 1; Dr. P Scambler for providing cosmids (sc11.1 and sc4.1), Dr. A Aurias for performing FISH in case 2, and Dr. MF Aillaud for performing the flow cytometry study of platelet glycoproteins. This work was supported by a grant from the Groupement de Recherches et d'Etudes sur les Génomes (GIP-GREG) and from EEC (Biomed 1).

REFERENCES

- Budarf ML, Konkle B, Michaud D, Ludlow LB, Michaud D, Mengrong L, Yamashiro DJ, McDonald-McGinn D, Zackai E, Driscoll D (1995): Identification of a patient with Bernard-Soulier syndrome and a deletion in the DiGeorge/velo-cardio-facial chromosome region in 22q11. *Hum Mol Genet* 4:763-766.
- D'Angelo JA, Pillers DM, Rice ML, Jett PL, Beyl CO, Hayflick S, Magenis SE (1994): Tetralogy of Fallot associated with deletion in the DiGeorge region of chromosome 22. *Am J Hum Genet* 55:A101.
- Driscoll DA, Salvin J, Sellinger B, Budarf M, McDonald-McGinn DM, Zackai, Emanuel BS (1993): Prevalence of 22q11 deletions in DiGeorge and velocardiofacial syndromes: Implications for genetic counselling and prenatal diagnosis. *J Med Genet* 30:913-917.



Fig. 4. Frontal view of family 2, mother and child. In the mother, the nose is typical with a large and prominent root.

- Goldmuntz E, Driscoll D, Budarf M, Zackai AH, McDonald-McGinn DM, Biegel J, Emanuel BS (1993): Microdeletion of chromosomal region in patients with congenital conotruncal cardiac defects. *J Med Genet* 30:807–812.
- Jyonouchi H, Voss RM, Krishna S, Urval K, Sjahli H, Welty PB, Good RA (1991): Soluble FcεR II in normal children and patients with immunodeficiency diseases. *J Allergy Clin Immunol* 87:965–970.
- Levy-Mozziconacci A, Wernert F, Scambler P, Rouault F, Metras D, Kreitman B, Depetris D, Mattei MG, Philip N (1994): Clinical and molecular study of DiGeorge sequence. *Eur J Pediatr* 153:813–820.
- Lipson A, Emanuel B, Colley P, Fagan K, Driscoll DA (1994): “CATCH 22” sans cardiac anomaly, thymic hypoplasia, cleft palate and hypocalcemia: Catch 22: A common result of 22q11 deficiency? *J Med Genet* 31:741.
- Nickel RE, Pillers DEA, Merckens M, Magen Re, Driscoll D, Emanuel B, Zonana J (1994): Velo-cardio-facial syndrome and DiGeorge sequence with myelomeningocele and deletions of the 22q11 region. *Am J Med Genet* 52:445–449.
- Sciré G, Dallapiccola B, Iannetti P, Bonaiuto F, Galasso C, Mingarelli R, Boscherini B (1994): Hypoparathyroidism as a major manifestation in two patients with 22q11 deletions. *Am J Med Genet* 52:478–482.
- Shprintzen RJ, Goldberg RB, Levin ML, Sidoti EJ, Berkman MD, Argamaso RV, Young D (1978): A new syndrome involving cleft palate, cardiac anomaly, typical facies, and learning disabilities: Velo-cardio-facial syndrome. *Cleft Palate J* 15:56–62.
- Shprintzen RJ, Goldberg R, Golding-Kushner KJ, Marion RW (1992): Late-onset psychosis in the velo-cardio-facial syndrome. *Am J Med Genet* 42:141–142.
- Wilson DI, Burn J, Scambler P, Goodship J (1993): DiGeorge syndrome: Part of CATCH 22. *J Med Genet* 30:852–856.